

TO KNOW ABOUT

MEDETOMIDINE

Risks With Medetomidine

- Slow breathing & heart rate
- In & out of awareness
- Increases the chance of overdose death
- Can cause severe withdrawal
 - Very fast heartbeat & high blood pressure
 - Heavy sweating, shaking, or twitching
 - Restlessness, anxiety, or panic

What is Medetomidine? ALSO KNOWN AS 'RHINO TRANQ'

- Often mixed with fentanyl
- Found in the street drug supply
- Used in animals and NOT approved for humans
- Stronger & longer-lasting than xylazine
- Causes deep sleep/sedation, slowed breathing, & low heart rate
- Two cases were reported in Tennessee emergency rooms in 2025, but it's unknown how common the drug is in the state





DETECTED IN TN DRUG SUPPLY



DANGEROUSLY SLOWS RREATHING

WHAT YOU CAN DO TO PROTECT YOURSELF FROM

Overdose Prevention

- Go slow & do not use alone
- Carry Naloxone & know how to use it
- Start rescue breathing right away
- Use drug checking strips to test for fentanyl and xylazine

Overdose Response

- □ Check for responsiveness
 - Shake & call out to the person
 - Perform a center chest rub
- □ Check for breathing
- □ Give them Naloxone every 2-3 mins
- □ Call 911 & stay with the person
- □ Start rescue breathing





MEDETOMIDINE

Resources



HOTLINE TO FIND
TREATMENT &
RECOVERY SERVICES



OVERDOSE RESPONSE
SAFETY HOTLINE
SAFE-SPOT.ME



HEALTH CRISIS
COUNSELING

SCAN FOR NALOXONE RESOURCES



TN'S SYRINGE SERVICE
PROGRAMS
TINYURL.COM/TNSSPS

